Department of Chemical and Physical Sciences Colloquium Seminar Series Wednesday, November 7, 2018 3:10pm in CC2150

DR. STEPHANIE WEBER DEPARTMENT OF BIOLOGY MCGILL UNIVERSITY

Organelles without borders: how liquid droplets organize the cell



Living cells contain a complex mixture of macromolecules. Typically, membranes partition these molecules into functional compartments called organelles. However, cells also contain compartments that lack an enclosing membrane and instead consist of local concentrations of protein and RNA. These compartments rapidly exchange components with the surrounding cytoplasm or nucleoplasm, raising a fundamental question: how do these compartments assemble and stably persist without a membrane holding them together? To answer this question, my lab uses quantitative live-cell imaging and physical modeling of the nucleolus, a prominent membraneless organelle responsible for ribosome biogenesis. In early C. elegans embryos, the bulk concentration of nucleolar components determines whether the nucleolus assembles and, if so, its size. These observations are consistent with a first-order phase transition, suggesting that nucleoli form by liquid-liquid phase separation of the nucleoplasm. Indeed, the coarsening dynamics of nucleoli are consistent with theoretical predictions of classical thermodynamic models, despite nonequilibrium activity in the cell. Our results suggest that phase separation represents a new principle of intracellular organization, capable of generating dynamic compartments of appropriate size.